



ANAPHYLAXIS POLICY

PURPOSE

To explain to Orchard Grove Primary School parents, carers, staff and students the processes and procedures in place to support students diagnosed as being at risk of suffering from anaphylaxis. This policy also ensures that Orchard Grove Primary School is compliant with Ministerial Order 706 and the Department of Education's guidelines for anaphylaxis management.

SCOPE

This policy applies to:

- all staff, including casual relief staff and volunteers
- all students who have been diagnosed with anaphylaxis, or who may require emergency treatment for an anaphylactic reaction, and their parents and carers.

POLICY

School Statement

Orchard Grove Primary School will fully comply with Ministerial Order 706 and the associated guidelines published by the Department of Education.

Anaphylaxis

Anaphylaxis is a severe allergic reaction that occurs after exposure to an allergen. The most common allergens for school-aged children are nuts, eggs, cow's milk, fish, shellfish, wheat, soy, sesame, latex, certain insect stings and medication.

Symptoms

Signs and symptoms of a mild to moderate allergic reaction can include:

- swelling of the lips, face and eyes
- hives or welts
- tingling/stinging in the mouth

Signs and symptoms of anaphylaxis, a severe allergic reaction, can include:

- difficult/noisy breathing
- swelling of tongue
- swelling/tightness in throat
- difficulty talking and/or hoarse voice
- wheeze or persistent cough
- persistent dizziness or collapse
- student appears pale or floppy
- abdominal pain and/or vomiting.

Symptoms usually develop within ten minutes and up to two hours after exposure to an allergen, but can appear within a few minutes.

Treatment

Adrenaline given as an injection into the muscle of the outer mid-thigh is the first aid treatment for anaphylaxis. Call an ambulance immediately following administration or concurrently if more than one adult is available at the scene.

Individuals diagnosed as being at risk of anaphylaxis are prescribed an adrenaline autoinjector for use in an emergency. These adrenaline autoinjectors are designed so that anyone can use them in an emergency.

Individual Anaphylaxis Management Plans

All students at Orchard Grove Primary School who are diagnosed by a medical practitioner as being at risk of suffering from an anaphylactic reaction must have an Individual Anaphylaxis Management Plan. When notified of an anaphylaxis diagnosis, the Principal of Orchard Grove Primary School will ensure that an Individual Anaphylactic Management Plan is developed, in consultation with the student's parents/carers.

Where necessary, an Individual Anaphylaxis Management Plan will be in place as soon as practicable after a student enrolls at Orchard Grove Primary School and where possible, before the student's first day.

Parents and carers must:

- obtain an ASCIA (Australasian Society of Clinical Immunology and Allergy) Action Plan for Anaphylaxis from the student's medical practitioner and provide a copy to the school as soon as practicable;
- immediately inform the school in writing if there is a relevant change in the student's medical condition and obtain an updated ASCIA Action Plan for Anaphylaxis;
- provide an up-to-date photo of the student for the ASCIA Action Plan for Anaphylaxis when that Plan is provided to the school and each time it is reviewed;
- provide the school with a current adrenaline autoinjector for the student that has not expired;
- participate in annual reviews of the student's Anaphylactic Management Plan.

Each student's Individual Anaphylaxis Management Plan must include:

- information about the student's medical condition that relates to allergies and the potential for anaphylactic reaction, including the type of allergies the student has;
- information about the signs or symptoms the student might exhibit in the event of an allergic reaction based on a written diagnosis from a medical practitioner;
- strategies to minimise the risk of exposure to known allergens while the student is under the care or supervision of school staff, including in the school yard, at camps and excursions, or at special events conducted, organised or attended by the school;

- the name of the person(s) responsible for implementing the risk minimisation strategies, which have been identified in the Plan;
- information about where the student's medication will be stored;
- the student's emergency contact details;
- an up-to-date ASCIA Action Plan for Anaphylaxis completed by the student's medical practitioner.

Review and updates to Individual Anaphylaxis Management Plans

A student's Individual Anaphylaxis Management Plan will be reviewed and updated on an annual basis in consultation with the student's parents/carers. The plan will also be reviewed and, where necessary, updated in the following circumstances:

- as soon as practicable after the student has an anaphylactic reaction at school;
- if the student's medical condition, insofar as it relates to allergy and the potential for anaphylactic reaction, changes;
- when the student is participating in an off-site activity, including camps and excursions, or at special events including fetes and concerts.

It is the responsibility of the parents/carers to inform the school if their child's medical condition changes.

Our school may also consider updating a student's Individual Anaphylaxis Management Plan if there is an identified and significant increase in the student's potential risk of exposure to allergens at school.

Location of plans and adrenaline autoinjectors

A copy of each student's Individual Anaphylaxis Management Plan will be stored with their ASCIA Action Plan for Anaphylaxis in the First Aid room, together with the student's adrenaline autoinjector. Adrenaline autoinjectors must be labelled with the student's name.

Adrenaline autoinjectors for general use are available at First Aid Room and are labelled "general use".

Risk Minimisation Strategies

To reduce the risk of a student suffering from an anaphylactic reaction at Orchard Grove Primary School, we have put in place the following strategies:

- staff and students are regularly reminded to wash their hands after eating;
- students are discouraged from sharing food;
- garbage bins at school are to remain covered with lids to reduce the risk of attracting insects;
- gloves must be worn when picking up papers or rubbish in the playground;
- if a school canteen is in use, staff are trained in appropriate food handling to reduce the risk of cross-contamination;
- year groups will be informed of allergens that must be avoided in advance of class parties, events or birthdays;
 - use non-food treats where possible, but if food treats are used in class it is recommended that parents of students with food allergy provide a clearly labelled treat box with alternative treats. Treats for other students should be handled with absolute care to ensure there is no cross contamination of food. If possible, foods should not contain the substances to which a student is allergic to.
- planning for off-site activities will include risk minimisation strategies for students at risk of anaphylaxis including supervision requirements, appropriate number of trained staff,

emergency response procedures and other risk controls appropriate to the activity and students attending;

- If a student does not have a current medically signed ASCIA plan and current EpiPen/Anapen, the school reserves that right to refuse the student to attend camp and/or excursions.

Adrenaline autoinjectors for general use

Orchard Grove Primary School will maintain a supply of adrenaline autoinjectors for general use, as a back-up to those provided by parents and carers for specific students, and also for students who may suffer from a first time reaction at school.

Adrenaline autoinjectors for general use will be stored in the first aid room and labelled “general use”.

The Principal is responsible for arranging the purchase of adrenaline autoinjectors for general use, and will consider:

- the number of students enrolled at Orchard Grove Primary School at risk of anaphylaxis
- the accessibility of adrenaline autoinjectors supplied by parents
- the availability of a sufficient supply of autoinjectors for general use in different locations at the school, as well as at camps, excursions and events
- the limited life span of adrenaline autoinjectors, and the need for general use adrenaline autoinjectors to be replaced when used or prior to expiry
- the weight of the students at risk of anaphylaxis to determine the correct dosage of adrenaline autoinjector/s to purchase.

Emergency Response

In the event of an anaphylactic reaction, the emergency response procedures in this policy must be followed, together with the school’s general first aid procedures, emergency response procedures and the student’s Individual Anaphylaxis Management Plan.

A complete and up-to-date list of students identified as being at risk of anaphylaxis is maintained by the First Aid Officer and stored at the first aid room. For camps, excursions and special events, a designated staff member will be responsible for maintaining a list of students at risk of anaphylaxis attending the special event, together with their Individual Anaphylaxis Management Plans and adrenaline autoinjectors, where appropriate.

If a student experiences an anaphylactic reaction at school or during a school activity, school staff must:

Step	Action
1.	<ul style="list-style-type: none">● Lay the person flat● Do not allow them to stand or walk● If breathing is difficult, allow them to sit ● Be calm and reassuring● Do not leave them alone● Seek assistance from another staff member or reliable student to locate the student’s adrenaline autoinjector or the school’s general use autoinjector, and the student’s Individual Anaphylaxis Management Plan, stored at first aid room● If the student’s plan is not immediately available, or they appear to be experiencing a first time reaction, follow steps 2 to 5
2.	<p>Administer an EpiPen or EpiPen Jr</p> <ul style="list-style-type: none">● Remove from plastic container

	<ul style="list-style-type: none"> ● Form a fist around the EpiPen and pull off the blue safety release (cap) ● Place orange end against the student’s outer mid-thigh (with or without clothing) ● Push down hard until a click is heard or felt and hold in place for 3 seconds ● Remove EpiPen ● Note the time the EpiPen is administered ● Retain the used EpiPen to be handed to ambulance paramedics along with the time of administration <p>OR</p> <p>Administer an Anapen® 500, Anapen® 300, or Anapen® Jr.</p> <ul style="list-style-type: none"> ● Pull off the black needle shield ● Pull off grey safety cap (from the red button) ● Place needle end firmly against the student's outer mid-thigh at 90 degrees (with or without clothing) ● Press red button so it clicks and hold for 10 seconds ● Remove Anapen® ● Note the time the Anapen is administered ● Retain the used Anapen to be handed to ambulance paramedics along with the time of administration
3.	Call an ambulance (000)
4.	If there is no improvement or severe symptoms progress (as described in the ASCIA Action Plan for Anaphylaxis), further adrenaline doses may be administered every five minutes, if other adrenaline autoinjectors are available.
5.	Contact the student’s emergency contacts.

If a student appears to be having a severe allergic reaction but has not been previously diagnosed with an allergy or being at risk of anaphylaxis, school staff should follow steps 2 – 5 as above.

Schools can use either the EpiPen® **and Anapen® on any student** suspected to be experiencing an anaphylactic reaction, regardless of the device prescribed in their ASCIA Action Plan.

Where possible, schools should consider using the correct dose of adrenaline autoinjector depending on the weight of the student. However, in an emergency if there is no other option available, any device should be administered to the student.

Note: If in doubt, it is better to use an adrenaline autoinjector than not to use it, even if in hindsight the reaction is not anaphylaxis. Under-treatment of anaphylaxis is more harmful and potentially life threatening than over-treatment of a mild to moderate allergic reaction. Refer to ‘Frequently asked questions’ on the [Resources tab](#) of the Department’s Anaphylaxis Policy.

Communication Plan

This policy will be available on Orchard Grove Primary School’s website so that parents and other members of the school community can easily access information about Orchard Grove Primary School’s anaphylaxis management procedures. The parents and carers of students who are enrolled at Orchard Grove Primary School and are identified as being at risk of anaphylaxis will also be provided with a copy of this policy.

The Principal is responsible for ensuring that all relevant staff, including casual relief staff, canteen staff and volunteers are aware of this policy and Orchard Grove Primary School's procedures for anaphylaxis management. Casual relief staff and volunteers who are responsible for the care and/or supervision of students who are identified as being at risk of anaphylaxis will also receive a verbal briefing on this policy, their role in responding to an anaphylactic reaction and where required, the identity of students at risk.

The Principal is also responsible for ensuring relevant staff are trained and briefed in anaphylaxis management, consistent with the Department's [Anaphylaxis Guidelines](#).

Staff training

The Principal will ensure that the following school staff are appropriately trained in anaphylaxis management:

- All staff are to complete the ASCIA Anaphylaxis e-training (<https://etrainingvic.allergy.org.au/login/index.php>) for Victorian Schools followed by a competency check in using an autoinjector tested in person within 30 days of completing the course. The e-training provided by ASCIA is valid for 2 years. A record of training and competency checks will be kept.
- OR
 - an approved face-to-face anaphylaxis management training course in the last three years (22300VIC First Aid Course Management of Anaphylaxis)
- A minimum of two staff members from the School will undertake face-to-face training to skill them in providing competency checks to assess their colleagues' ability to use an EpiPen and become School Anaphylaxis Supervisors. This is: Course in Verifying the Correct Use of Adrenaline Autoinjector Devices 22303VIC. (valid for 3 years)
- New staff will be required to complete training as soon as practicable.

Staff are also required to attend a briefing on anaphylaxis management and this policy at least twice per year (with the first briefing to be held at the beginning of the school year), facilitated by a staff member who has successfully completed an anaphylaxis management course within the last 3 years. Each briefing will address:

- this policy
- the causes, symptoms and treatment of anaphylaxis
- the identities of students with a medical condition that relates to allergies and the potential for anaphylactic reaction, and where their medication is located
- how to use an adrenaline autoinjector, including hands-on practice with a trainer adrenaline autoinjector
- the school's general first aid and emergency response procedures
- the location of, and access to, adrenaline autoinjectors that have been provided by parents or purchased by the school for general use.

When a new student enrolls at Orchard Grove Primary School who is at risk of anaphylaxis, the Principal will develop an interim plan in consultation with the student's parents and ensure that appropriate staff are trained and briefed as soon as possible.

A record of staff training courses and briefings will be maintained in Google docs and through the school's online Emergency Management Plan.

The Principal will ensure that while students at risk of anaphylaxis are under the care or supervision of the school outside of normal class activities, including in the school yard, at camps and excursions,

or at special event days, there is a sufficient number of school staff present who have been trained in anaphylaxis management.

FURTHER INFORMATION AND RESOURCES

- The Department's Policy and Advisory Library (PAL):
 - [Anaphylaxis](#)
- [Allergy & Anaphylaxis Australia](#)
- ASCIA Guidelines: [Schooling and childcare](#)
- Royal Children's Hospital: [Allergy and immunology](#)
-

The following school policies are also relevant to this Anaphylaxis Policy:

- Administration of Medication
- First Aid
- Asthma
- Health Care Needs

Review Cycle And Evaluation

Ratified May 2023
Schedule for review in 2024

The Principal will complete the Department's Annual Risk Management Checklist for anaphylaxis management to assist with the evaluation and review of this policy and the support provided to students at risk of anaphylaxis.

APPENDIX 1:

Raising Student Awareness

Peer support is an important element of support for students at risk of anaphylaxis. Staff can raise awareness in school through fact sheets or posters displayed in hallways and classrooms.

Class teachers will discuss the topic with students in class, with a few simple key messages such as:

Student messages about anaphylaxis	
1.	Always take food allergies seriously - severe allergies are no joke.
2.	Don't share your food with friends who have food allergies.
3.	Wash your hands after eating.
4.	Know what your friends are allergic to.
5.	If a school friend becomes sick, get help immediately.
6.	Be respectful of a school friend's EpiPen.
7.	Don't pressure your friends to eat food that they are allergic to.

Source: Be a Mate kit, published by Anaphylaxis and Allergy Australia

Related documents

- Enrolment checklist
- ASCIA action plan
- Individual Management Plan
- Brochure titled "Anaphylaxis-a life threatening reaction", available through the Royal Children's Hospital, Department of Allergy

Contact details for resources and support

Royal Children's Hospital, Anaphylaxis Advisory Line -1300 725 911 (toll free) or 9345 4235

Anaphylaxis Australia

- www.allergyfacts.org.au 1300 728 000 ASCIA

- <http://www.allergy.org.au/schools-childcare>

APPENDIX 2:

OGPS Anaphylaxis Management Plan

This Plan is to be completed by the principal or nominee on the basis of information from the student's medical practitioner provided by the parent/carer.

APPENDIX 3:

The following procedures are to be put in place in the case of suspected anaphylactic reaction.

If an allergen turns up in the classroom


1	Remove the allergen and the carrier of the allergen from the immediate environment
2	Support the anaphylactic student by ensuring they are removed from the site of the allergen (preferably in the same room- we do not want to cause panic in the student) with an adult or student with them to monitor and watch for any possible reaction
3	Contact the office for First Aid Officer to attend
4	Area of contamination to be immediately cleaned (eg. tables are to be wiped down)
5	First Aid Officer to go immediately to area of contamination to support the anaphylactic student and the staff member/volunteer/relief teacher on duty
6	First Aid Officer/Office is to contact any families directly involved
7	At the Principal's discretion, communication may be sent home to all families from the group of students regarding anaphylaxis awareness at OGPS.

APPENDIX 4:


How to Administer an EpiPen or EpiPen Jr:

Remove EpiPen from plastic container.


How to give EpiPen® or EpiPen® Jr adrenaline (epinephrine) autoinjectors



1. Form fist around EpiPen® and PULL OFF BLUE SAFETY RELEASE.



2. Hold leg still and PLACE ORANGE END against outer mid-thigh (with or without clothing).



3. PUSH DOWN HARD until a click is heard or felt and hold for 3 seconds

REMOVE EpiPen®

Note: All EpiPen®s should be held in place for 3 seconds regardless of instructions on device label


Note the time EpiPen administered (record on hand of the side EpiPen administered).
Return EpiPen to its plastic container.

In the rare situation where there is no marked improvement and severe symptoms are present, a second injection may be administered after 5 minutes, if available.

How to give Anapen® adrenaline (epinephrine) autoinjectors




PULL OFF BLACK
NEEDLE SHIELD



PULL OFF GREY
SAFETY CAP
from red button



PLACE NEEDLE END
FIRMLY against outer
mid-thigh at 90° angle
(with or without clothing)



PRESS RED BUTTON
so it clicks and hold
for 10 seconds.
REMOVE Anapen®

After Anapen® is used, put it to one side and do not touch the exposed needle.

After phoning ambulance, place needle into wide end of the black needle shield, or place the used Anapen® in a container. Provide ambulance with the used Anapen® and the time it was given.